

| | |
|-------|-------------|
| レジメン名 | Bev+TAS-102 |
| がん種 | 大腸癌 |

| | | | |
|-------------|-------------------------|----|--|
| 一般名 | 投与量 | 単位 | |
| ベバシズマブ | 5 mg/kg | | |
| TAS-102(内服) | 70 mg/m ² /日 | | |




4週で1クール

5日間服薬 → 休 → 5日間服薬 → 休 → 2週間休薬

day 1 ... 8 ... 15 28

1クール

TAS-102は、Day1夕より、5日間服薬・2日間休薬を2回繰り返した後、2週間休薬する。

| | 治療内容 | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|--|--|----|---|-----|---|---|---|-----|---|---|---|-----|---|---|---|-----|---|---|---|-----|---|---|---|-----|---|---|---|--|
| | メイン | 側管 | | | | | | | | | | | | | | | | | | | | | | | | | | |
| <p>ルート確保用輸液は、所要時間に合わせて適宜速度調節する。治療薬の投与中は中止可。残破棄可。</p> <div style="border: 1px solid black; padding: 5px; margin: 5px 0;">生食(100ml)</div> <div style="border: 2px solid red; border-radius: 15px; padding: 10px; margin: 5px 0;"> <p>⚠ 注意 ベバシズマブの投与時間は、初回90分。 投与に問題がなければ、次回より60分→30分と短縮可能</p> </div> <div style="border: 1px solid gray; border-radius: 15px; padding: 10px; margin: 5px 0;"> <p>ロンサーフは、朝、夕食後に服用。 day1夕からday6朝まで、および、day8夕からday13朝まで服用。 !! 服用量を確認 !!</p> <div style="display: flex; justify-content: space-around;">   </div> </div> | <p>経過時間</p> <table style="width: 100%; border-collapse: collapse;"> <tr><td style="border: 1px solid black;">0</td><td style="border: 1px solid black;">←</td></tr> <tr><td style="border: 1px solid black;">:30</td><td style="border: 1px solid black;">↓</td></tr> <tr><td style="border: 1px solid black;">1</td><td style="border: 1px solid black;">↓</td></tr> <tr><td style="border: 1px solid black;">:30</td><td style="border: 1px solid black;">↓</td></tr> <tr><td style="border: 1px solid black;">2</td><td style="border: 1px solid black;">←</td></tr> <tr><td style="border: 1px solid black;">:30</td><td style="border: 1px solid black;">↓</td></tr> <tr><td style="border: 1px solid black;">3</td><td style="border: 1px solid black;">↓</td></tr> <tr><td style="border: 1px solid black;">:30</td><td style="border: 1px solid black;">↓</td></tr> <tr><td style="border: 1px solid black;">4</td><td style="border: 1px solid black;">↓</td></tr> <tr><td style="border: 1px solid black;">:30</td><td style="border: 1px solid black;">↓</td></tr> <tr><td style="border: 1px solid black;">5</td><td style="border: 1px solid black;">↓</td></tr> <tr><td style="border: 1px solid black;">:30</td><td style="border: 1px solid black;">↓</td></tr> <tr><td style="border: 1px solid black;">6</td><td style="border: 1px solid black;">↓</td></tr> </table> | 0 | ← | :30 | ↓ | 1 | ↓ | :30 | ↓ | 2 | ← | :30 | ↓ | 3 | ↓ | :30 | ↓ | 4 | ↓ | :30 | ↓ | 5 | ↓ | :30 | ↓ | 6 | ↓ | <div style="border: 1px solid black; padding: 5px; margin: 5px 0;">ベバシズマブBS 生食(100ml)</div> <div style="border: 1px solid black; padding: 5px; margin: 5px 0;">生食(50ml) フラッシュ用</div> <div style="border: 1px solid gray; border-radius: 15px; padding: 10px; margin: 5px 0;"> <p>針先が、抗癌剤を含まない輸液に満たされた後、抜針する</p>  </div> |
| 0 | ← | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| :30 | ↓ | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 1 | ↓ | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| :30 | ↓ | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 2 | ← | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| :30 | ↓ | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 3 | ↓ | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| :30 | ↓ | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 4 | ↓ | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| :30 | ↓ | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 5 | ↓ | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| :30 | ↓ | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 6 | ↓ | | | | | | | | | | | | | | | | | | | | | | | | | | | |